

SURFICIAL GEOLOGY OF THE ALTAMONT 7.5-MINUTE QUADRANGLE, ALBANY AND SCHENECTADY COUNTIES, NEW YORK

prepared by
Sean P. Grasing and Karl J. Backhaus

Supported in part by the U.S Geological Survey's
National Cooperative Geologic Mapping Program STATEMAP Award Number G16AC00293

INTRODUCTION:

The Altamont 7.5-Minute Quadrangle was mapped as part of the National Cooperative Geologic Mapping Program's Great Lakes Geologic Mapping Coalition projects in 2015, supported by award #G16AC00293. This quadrangle is one of 18 that have been mapped under the Albany County Surficial Geologic Mapping Project, which the NYSGS initiated in 2015 and aims to conclude in the early 2020s. The purpose of this map is to identify and delineate various surficial and geologic materials with the intent to inform and guide municipalities in land-use, environmental, and natural resource decisions across its approximately 54-square-mile area.

The mapping area is situated between the Catskill Mountains and Hudson-Mohawk Valley Lowlands in New York State. The Altamont Quadrangle extends across two counties encompassing the southern border of Schenectady County and the northern border of Albany County. The Village of Altamont, along with the towns of Berne, Knox, and New Scotland are the primary municipalities in the Altamont quadrangle. John Boyd Thacher and Thompsons Lake State Parks, along with the Winn and Wolf Hill Preserves, are located within the quadrangle.

The Altamont Quadrangle is a part of the Catskill Mountains and Hudson-Mohawk Valley Lowlands physiographic provinces and features an escarpment as well as lowlands. Below the escarpment, one can find glacial lake deposits that were formed during the period of Lake Albany, along with some sand deposits that were formed afterwards under aeolian conditions. The uplands region encompasses the Helderberg Escarpment, a bedrock escarpment that has a slight south-southwest dip (Ver Straeten and others, 2013). This escarpment extends diagonally across the quadrangle, from the southwestern corner to the northwest corner. The elevation within the entire quadrangle ranges widely, with the lowest elevations lying at 263 feet above mean sea level (feet-amsl; 86 meters-amsl) near West Old State Road and the highest elevation at 1,819 feet-amsl (554 meters-amsl) on top of Mount Helderberg. Prominent bodies of water in the quadrangle are Thompsons Lake, Duane Lake, Warner Lake, and the Altamont Reservoir. The Bozenkill Creek flows through the middle of the quadrangle.

The bedrock of the Altamont Quadrangle is composed solely of sedimentary rocks, including limestone, shale, mudstone, and sandstone (Ver Straeten and others, 2013). These rocks range in age from Late Ordovician to Late Devonian. Until it reaches the escarpment, the Hudson-Mohawk Valley in the lowlands of the quadrangle is primarily characterized by the Late Ordovician Schenectady Formation. At the escarpment the valley contacts the Helderberg Group, which includes the Marlius, Rondout, Coeymans, Kalkberg, New Scotland, and Becraft Formations. On the top of the escarpment there are outcrops of the Otiskany, Esopus, Schoharie, Onondaga, and Union Springs Formations, as well as the Marcellus Subgroup.

The surficial geologic units in this quadrangle were previously mapped at a scale of 1:250,000 and were identified as outwash sand and gravel, swamp deposits, alluvium, kame, till, rock, and lacustrine sand and deltas (Cadwell et al., 1991).

METHODOLOGY:

Preliminary field maps of the Altamont Quadrangle were generated as an initial step using ESRI ArcMap software. These maps incorporated all available topographic data, including roads, lidar surface terrain, and hydrography, and served as a base for plotting field data. Field data comprised field stops, bedrock outcrops, and other important site information. A five-and-a-half-foot hand auger was used to enable surficial sampling below the variably thick organic soil horizon, which is situated beneath the topsoil. An entrenching shovel and pick were also used to remove topsoil and/or eroded sediments from outcrops or exposures, thereby exposing fresh sediment for analysis. At each field stop, meticulous details were logged into a field notebook including: coordinates captured by a Garmin GPS 550i, descriptive notes on the sediment encountered, the names of samples taken, the time of the stop, and any high-resolution photographs taken.

At most of the field sampling sites, soil samples were collected for grain-size analysis. This involved either dry-sieve or wet-sieve analysis. Both methods followed the procedure outlined by Bowles (1978), but utilized a seven-tiered sieve stack (#5, #10, #18, #35, #60, #120, #230, and Pan) for both dry (mechanical) and wet (hydrometer) sieve analysis. Predominantly cohesive samples, which are fine-grain dominant, were analyzed using wet-sieve methods, while cohesionless samples, which are coarse-grain dominant, were analyzed using dry-sieve methods.

The final surficial geologic map, along with cross sections and elevation maps, was generated using ESRI ArcMap and Adobe Illustrator CS6 software. Cross sections were initially created in ArcMap utilizing the XActo Cross Section 10 Tool. These cross sections were then exported to Adobe Illustrator for the correlation of stratigraphic units. The surficial geologic map was developed by scanning mylar sheets that had been drafted based on the geologic field maps. Polygons were then created by digitizing these scanned maps in ArcMap and color-coding them to represent the surficial geologic units within the quadrangle. The final map was assembled in Adobe Illustrator and exported as a PDF file.

RESULTS:

A total of 141 field stops were conducted in the quadrangle, from which 100 samples were collected for grain-size analysis. Multiple samples were obtained from stops where stratigraphy was observed either on the surface or at depths that could be reached using a hand auger. The lithologies discovered during field sampling included: 97 were diamicton, exposed 35 were bedrock, four were sand, three were glaciolacustrine silt and clay and two were sand and gravel. The surficial geologic units identified within the quadrangle are as follows:

Artificial Fill (Af)

This unit generally consists of coarse-to-fine materials such as large cement mounds and crushed rock, which have been transported anthropogenically for construction purposes. These materials are commonly used in large artificial dams designed to retain water, elevated roads, bridge abutments, quarries and large mill facilities within the quadrangle that are raised above the surface.

Holocene Alluvium (Ha) and Holocene Wetland Deposits (Hw)

Post-glacial sediments are found in low-lying areas or land depressions throughout the quadrangle. Holocene alluvium is associated with fluvial processes in creek valleys across the area and primarily silt, sand, and stratified silt, sand, and gravel. Holocene wetland deposits occur in low areas and depressions in the highlands of the quadrangle, where wetlands form due to limited drainage capacity. The lithology in these areas generally comprises peat, marl, clay, or sand.

Diamict Colluvium (Hdc)

Unsorted and unstratified deposit of gravel, sand, silt, clay, with boulders/cobbles possible. Described as a mass-wasting deposit at the base of steep hillslopes and cliffs as part of a slump or hillslope failure. Located in minuscule-sized deposits along Bozenkill Creek and large slopes along the Helderberg Escarpment.

Pleistocene Silt and Clay (Pisc)

Stratified, fine-grained sediment consisting of fine sand, silt and clay size particles. Inferred to be deposited in mid-shore to deepwater settings of glacial lakes. May include marl, rhythmites, and varves. Located in the lowlands of Altamont quadrangle.

Pleistocene Sand (Ps)

Well-sorted and stratified sand is deposited through fluvial, lacustrine, or aeolian processes. These are inferred to be deposits associated with distal glacial environments. Observations indicate that these well-sorted sand deposits often overlie coarser sand and gravel deposits (Psg), likely because of a decrease in depositional energy or potential aeolian activity. Located in the northeastern corner of the Altamont quadrangle, along with loess deposits.

Pleistocene Sand and Gravel (Psg)

Characterized by well-sorted and stratified sand and gravel, this unit is believed to have been deposited by glacial meltwater at or very near the glacier's edge. It is often found at elevations several meters higher than the floors of present-day river valleys. Located in the lowlands of Altamont quadrangle.

Pleistocene Cobbles to Sand (Pics)

Stratified ice-contact deposits, variable coarse-grained sediment consisting of boulders to sand size particles. Inferred to be deposited with stagnant ice in the form of sand and gravel hummocks. Found in the lowlands of the Altamont quadrangle.

Pleistocene Diamicton (Pd)

This unit consists of a heterogeneous mixture of sediment grains, ranging in size from clay to boulders. In this quadrangle all diamicton is interpreted to be glacial till, which is sediment deposited directly beneath a glacier. The diamictons are predominantly located on top of the escarpment, where they are mostly clast-supported, and can form rather thick deposits here and in the Bozenkill Creek Valley.









SUMMARY AND CONCLUSIONS:

The surficial geology in the Altamont Quadrangle is characterized by five distinct informal regions: the lowlands, Settles Hill, the Bozenkill Creek Valley, the Helderberg Escarpment, and the uplands.


The lowlands define the region east of the Helderberg Escarpment. Starting from Altamont, this zone travels south on NY 156 and north on NY 397 and is located to the east of either route. Starting from Altamont, this zone travels south on NY 156 and north on NY 397 and is located to the east of either route. These sediments are matrix-supported with no visible clasts, often mottled (red- and gray-colored spots) or gray, dry and stiff. Loess and sand (Ps) deposits are located on West Old State Road in the north-easternmost corner of the quadrangle. These deposits are well-sorted, either sand or silt dominated, and yellowish-brown. Sand and gravel (Psg) deposits are found around Gardner Road. They are matrix-supported and include pebbles and cobbles that are sub-rounded to well-rounded. Diamicton (Pd) deposits are commonplace as well in the lowlands. Their characteristics vary but are all poorly sorted. Pebbles and boulders within are typically very angular to rounded. North of NY 156, Gardner Road passes over ice-contact sand and gravel deposits (Pics) which consist of poorly sorted, sub-rounded to rounded sand and gravel with cobbles.

Settles Hill is north of the Bozenkill Creek, which Settles Hill Road and Furber Road follow. The hill has only diamicton (Pd) deposits. The southern portion of the hill has thicker deposits of diamicton while the entire northern section of the hill has shallow deposits even as you move towards the west. Although there are shallow deposits on Gray Road, which runs to the south. This suggests that the valley was eroded and then filled in the past. On top of Settles Hill there are a few drumlins and streamlined features that have a bearing ranging from 240 to 260 degrees. This suggests that the glacier flowed toward the Mohawk Valley and potentially belonged to either the Mohawk or Janesville-Cobleskill ice streams (Kozlowski, Backhaus, Frieman, Leone, and Ferenc, 2023). Settles Hill has different flow directions compared to the drumlins and streamlined features in the upland section of the quadrangle's southern region.

SYMBOLS

— Street	 Water Body	• NYSGS Soil Sample Location
 Highway	 Stream	+ NYDEC Water Well Location
 Railroad	 Contour	+ NYSDOT Boring Location
 County Line	 Cross-Section Line	 Drumlin



QUADRANGLE LOCATION



ADJOINING QUADRANGLES

Duanesburg	Rothsden Junction	Schenectady
Gallipolis	Altamont	Northville
Rensselaerville	Westerlo	Catskillville

QUADRANGLE ELEVATION



1:75,000 scale; 2x vertical exaggeration
Shaded relief generated from 2019 FEMA 1-meter lidar data set.

SURFICIAL GEOLOGY OF THE ALTAMONT 7.5-MINUTE QUADRANGLE, ALBANY AND SCHENECTADY COUNTIES, NEW YORK

Sean P. Grasing, James R. Leone and Andrew L. Kozlowski

2024

NOTICE

This geologic map was funded in part by the USGS National Cooperative Geologic Mapping Program STATEMAP award number G16AC00293 in the year 2016 & 2023.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily presenting the official policies, either expressed or implied, of the U.S. Government.

While every effort has been made to ensure the integrity of this digital map and the factual data upon which it is based, the New York State Education Department ("NYSED") makes no representation or warranty, expressed or implied, with respect to its accuracy, completeness, or usefulness for any particular purpose or scale. NYSED assumes no liability for damages resulting from the use of any information, apparatus, method, or process disclosed in this map and text, and urges independent site-specific verification of the information contained herein. Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by NYSED.

New York State Museum Map & Chart No. 232

ISSN:0097-3793 ; ISBN:978-1-55557-430-7